




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07C 69/732, 67/317, C12P 7/62	A1	(11) International Publication Number: WO 97/06128 (43) International Publication Date: 20 February 1997 (20.02.97)
(21) International Application Number: PCT/EP96/03495 (22) International Filing Date: 5 August 1996 (05.08.96) (30) Priority Data: 95202125.1 3 August 1995 (03.08.95) EP (34) Countries for which the regional or international application was filed: NL et al. 60/001,860 3 August 1995 (03.08.95) US (71) Applicant (for all designated States except US): GIST-BROCADES B.V. [NL/NL]; Waterringseweg 1, P.O. Box 1, NL-2600 MA Delft (NL). (72) Inventors; and (75) Inventors/Applicants (for US only): DE PATER, Robertus, Mattheus [NL/NL]; Verkadestraat 6, NL-2613 ZL Delft (NL). SIBEYN, Mieke [NL/NL]; Pauwstraat 8, NL-3816 AV Amersfoort (NL). (74) Agents: VISSER-LUIRINK, Gesina et al.; Gist-Brocades N.V., Patents and Trademarks Dept., Wateringseweg 1, P.O. Box 1, NL-2600 MA Delft (NL).	(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>	

(54) Title: SELECTIVE PROCESS FOR THE DEACYLATION OF ACYLATED COMPOUNDS

**(57) Abstract**

The invention provides a simple and selective method for the deacylation of 4-acylated statins during the preparation of statins from a fermentation broth, more specifically a reduction of impurities from the process, by increasing the pH of the fermentation broth.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

5 **SELECTIVE PROCESS FOR THE DEACYLATION OF ACYLATED COMPOUNDS****FIELD AND BACKGROUND OF THE INVENTION**

The present invention provides an improvement for the recovery of lovastatin, compactin or pravastatin from
10 fermentation broths.

Lovastatin for instance is produced as a secondary metabolite by various microorganisms such as Aspergillus terreus (US 4231938) or Monascus ruber (US 4323648). During the fermentation also lovastatin related byproducts such as 4-acetyl
15 lovastatin are produced.

Lovastatin, usually in the acid form, can be isolated from the fermentation broth in different ways. The first stage is formed by purification yielding crude crystals. These crude crystals still comprise related compounds like 4-acetyl
20 lovastatin. As lovastatin is a pharmaceutical compound that has to meet high purity requirements, additional purification in order to remove the lovastatin related impurities is necessary. The lovastatin-related impurities are generally removed by multiple recrystallizations, by column chromatography as
25 described in US patent 4231938 or preparative HPLC (WO 92/16276), decreasing the yield significantly.

By the process of the present invention impurities present in the broth filtrate are removed, thus preventing the need for
30 their removal via additional recrystallizations and resulting in an increased yield.

During the application of the process of the present invention to a broth filtrate of a microorganism producing statins as for instance Aspergillus terreus, surprisingly 4-acetyl lovastatin is selectively converted into lovastatin instead of being converted into dehydro lovastatin via dehydration which occurs for pure 4-acetyl lovastatin (see Figure I). Another surprising fact is that the 2-methyl

- 2 -

butanoate group is not removed during the application of the invention.

The process of the present invention has neither been described nor suggested in the prior art.

5

BRIEF DESCRIPTION OF THE FIGURES

I. Reaction scheme of the deacylation of 4-acetyl lovastatinic acid to lovastatinic acid (a) and the dehydration of 4-acetyl lovastatinic acid to dehydro lovastatinic acid (b)

II. Thin layer chromatogram (TLC) showing reduction of impurities in crude lovastatin crystals after applying the process of the invention. Eluent: chloroform/methanol=9/1; detection: iodine staining; run product: 2 μ l of a solution, consisting of crude lovastatin crystal in toluene, concentration 50 g/l.

Right side: crude crystal from untreated broth filtrate; left side: crude crystal from broth filtrate which has been stirred for 2 hours at 50°C and pH 12.5.

DESCRIPTION OF THE INVENTION

25

The present invention concerns a process to improve the recovery of lovastatin, pravastatin or compactin from a broth filtrate. This process comprises:

- growing in a medium microorganisms that produce a member of the group consisting of lovastatin, compactin or pravastatin resulting in a product medium
- removal of the biomass from the product medium to obtain a clarified broth filtrate
- isolating the purified lovastatin, compactin or pravastatin, respectively,

characterized by adjusting the pH of the clarified broth filtrate above about pH 10. Preferably, said method also

comprises heating the clarified broth filtrate above approximately 50°C.

The process of the present invention offers a simple and selective method of the deacylation of 4-acylated statins in
5 broth filtrates, resulting in an improved yield and purity of the crystals. During the treatment at high pH the 4-acylated statin is converted into the related statin. The conversion rate of, for instance, 4-acetyl lovastatin into lovastatin in broth filtrate is dependent on the pH and the reaction temperature.
10 In a preferred embodiment of the present invention, the treatment is carried out at pH-values above pH=10, more preferably between pH=10 and pH=13, most preferably between pH=11 and pH=12.5. Also temperatures between 60°C and 95°C are preferred. By applying higher pH-values and/or higher
15 temperatures the reaction time for complete deacylation decreases.

The process of increasing the pH can be advantageously be applied to filtrates of fermentation broths from any microorganism that is capable of producing a member of the group
20 consisting of lovastatin, pravastatin or compactin. Microorganisms capable of producing statins may be one of the following species:

Penicillium, Hypomyces, Paecilomyces, Eupenicillium,
Trichoderma, Aspergillus, Monascus, Phoma, Doratomyces,
25 Gymnoascus or Pleurotus.

The fermentation of these microorganisms in order to produce statins is carried out in aqueous media similar to those employed for the production of other fermentation products. Such media contain sources of carbon, nitrogen and inorganic salts
30 assimilable by the microorganism.

In general, carbohydrates such as sugars, for example glucose, maltose, sucrose, xylose, mannitol and the like and starches such as grains, for example, oats, ryes, cornstarch, corn meal and the like can be used either alone or in
35 combination as sources of assimilable carbon in the nutrient medium. These carbon sources can be used individually, or several such carbon sources may be combined in the medium.

In general, many proteinaceous materials may be used as nitrogen sources in the fermentation process. Suitable nitrogen sources include for example, yeast hydrolysates, primary yeast, yeast extracts, soybean meal, cottonseed flour, hydrolysates of
5 casein, corn steep liquor, distiller's solubles or tomato paste and the like. The sources of nitrogen can be used either alone or in combination.

Among the nutrient inorganic salts which can be incorporated in the culture media are the customary salts
10 capable of yielding sodium, potassium, ammonium, calcium, phosphate, sulfate, chloride, carbonate and like ions. Also included are trace metals such as cobalt, manganese, iron and magnesium.

It should be noted that the media described in the Examples
15 are merely illustrative of the wide variety of media which may be employed, and yet are not intended to be limitative. Specifically, the carbon sources used in the culture media to produce lovastatin included dextrose, dextrin, glucose, sucrose, oat flour, oatmeal, molasses, citrate, acetate, soybean oil,
20 glycerol, malt extract, cod liver oil, starch, ethanol, figs, ascorbate, and lard oil. Included as nitrogen sources were peptonized milk, autolyzed yeast, yeast extract, yeast RNA, tomato paste, casein, primary yeast, peanut meal, distillers solubles, corn steep liquor, soybean meal, corn meal, NZ amine,
25 beef extract, asparagine, cottonseed meal, ammonia and ammonium sulphate. The major ionic components, CaCO_3 , KH_2PO_4 , $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ and NaCl can also be added as well as small amounts of $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ and traces of Fe, Mn, Mo, B and Cu. The nutrients can either be dosed in the batch medium or can be (partly) fed during the
30 fermentation.

The process of the present invention is applied directly to the broth after removal of the biomass, prior to further purification steps. For instance, an aqueous solution of earth alkali hydroxide or ammonium hydroxide can be conveniently used
35 for such a reaction. The broth filtrate is treated at high pH and preferably at high temperatures. Furthermore, during this treatment at high pH, proteins present in the fermentation broth

filtrate are denaturated, facilitating their removal from the product in the subsequent reaction steps. Further purification steps may comprise extraction, adsorption to a hydrophobic resin, ionexchange, column chromatography etc.

5 The following examples will illustrate the invention and are offered by way of illustration and not by way of limitation. Experiments II to IV are carried out under a nitrogen atmosphere.

10

EXAMPLES

EXAMPLE I: PRODUCTION OF LOVASTATIN BY MEANS OF FERMENTATION OF ASPERGILLUS TERREUS STRAIN AD43

15 Aspergillus terreus strain AD43, DS number 28373 has been deposited with the Centraal Bureau voor Schimmelcultures (CBS, Delft, The Netherlands), and has been granted CBS accession number CBS 456.95.

20 One 1 ml vial of a spore suspension of Aspergillus terreus strain AD43, stored in glycerol at -80°C was opened aseptically, and contents were suspended in a 2 liter shake flask containing 500 ml of the following medium (heated in an autoclave for 20 minutes at 121°C):

25	Ingredient	Amount per kg
	Glucose.1H ₂ O	10 g
	Oatmeal	10 g
	Tomato paste	40 g
	Corn steep solids	5 g
30	Trace elements	1 g

Composition of the trace element solution (per 100 ml of distilled water): FeSO₄.7H₂O, 1 g; MnSO₄.1H₂O, 1 g; CuCl₂.2H₂O, 0.025 g; CaCl₂.2H₂O, 0.1 g; H₃BO₄, 0.056 g; (NH₄)₆Mo₇O₂₄.4H₂O, 35 0.019 g; ZnSO₄.7H₂O, 0.2 g.

The shake flask was incubated at 28°C during 24 hours in a rotary shaker at 280 rpm (throw of 3.5 cm). 20 ml of the shake

- 6 -

flask broth (diluted in 100 ml of saline solution) was then inoculated into a fermenter with 10 kg of broth weight. The composition of the fermentation broth was as follows:

5	Ingredient	Amount per kg
	Glucose.1H ₂ O	20 g
	Yeast extract paste	33 g
	Polypropylene glycol 2000	2.5 ml

10 Glucose and the yeast extract/polypropylene glycol solution were sterilized separately (20 minutes at 121°C).

 Fermentation conditions were as follows:

 pH was kept constant at 6.5, using H₂SO₄ and NaOH

 Temperature was 28°C

15 Air supply was 1 vvm

 As soon as all glucose was consumed a glucose/yeast extract feed was started at a rate of 1.2 g of glucose per kg of broth per hour. Composition of the feed:

20	Ingredient	Amount per kg
	Glucose.1H ₂ O	500 g
	Yeast extract paste	17 g
	Polypropylene glycol 2000	14 ml

25

 After 192 hours of fermentation the pH of the broth was raised to pH 10 with NaOH and the broth was diluted with 4 liters of water.

 This fermentation yielded 385 mg of lovastatin acid per
30 liter of fermentation broth before dilution. After dilution a lovastatin acid content of 411 mg/l was measured.

EXAMPLE II: EFFECT OF HEAT TREATMENT OF BROTH FILTRATE ON THE PURITY OF LOVASTATIN CRYSTALS

35

 1,000 ml of broth filtrate of strain AD43 (lovastatin acid concentration 0.4 g/l, produced according to example I) was

brought to pH 12.5 with 2 N NaOH at 25°C, and subsequently brought to 50°C for 2 hrs. After 2 hrs, the reaction was completed, and the reaction mixture was cooled to room temperature. Then the pH was lowered to pH 4 using sulfuric acid, 3,000 ml of toluene were added en mixed during 30 minutes.

The toluene layer was separated from the water layer, and subsequently concentrated to a volume of 80 ml by evaporation at 40°C under vacuum.

The lovastatin acid in the extract was converted into the lactone by heating it to 90°C for 3 hours (yield of conversion was 99.2%). After cooling to room temperature, the toluene was mixed with 80 ml of water, while the pH was adjusted to pH=10 with NaOH. After separation of the layers, the toluene layer was mixed again with 80 ml of fresh water, while the pH was adjusted to pH=4 with sulfuric acid. After separation of the layers, the toluene layer was treated with 0.1 g of active coal, Norit SX ultra. Subsequently the toluene solution was filtrated and further concentrated to 15 ml by evaporation. Cooling to -10°C resulted in crystallization. The crystals were washed with 5 ml of cold toluene, and dried under vacuum at room temperature. In these crystals no 4-acetyl lovastatin could be detected, neither by TLC nor by proton NMR.

In contrast, crystals obtained from the fermentation broth via the process described above but without a heat treatment of the broth filtrate at pH 12, did contain 4-acetyl lovastatin as detected by TLC (see Figure II). Also proton NMR-analysis showed the presence of 1.1% of 4-acetyl lovastatin in these crystals.

EXAMPLE III: COMPARISON OF VARIOUS HEAT TREATMENT CONDITIONS OF BROTH FILTRATE ON THE PURITY OF LOVASTATIN CRYSTALS

Various portions of broth filtrate were treated at different pH-values and temperatures and of different duration, as shown in Table 1. For each set of parameters, 1,000 ml of a broth filtrate of strain AD43 (lovastatin acid concentration of 0.4 g/l) was used. After the treatment, the filtrate was brought to pH=4 using sulfuric acid, and 1,000 ml of toluene were mixed

with the filtrate for 30 minutes. The layers were subsequently separated, and the toluene layer was concentrated to a volume of 80 ml by evaporation, and kept at 90°C for 3 hours. After cooling to room temperature, the toluene was mixed with 40 ml of water, while the pH was adjusted to pH=10 with NaOH. After separation of the layers, the toluene was mixed with another 40 ml of water, while the pH was adjusted to pH=4 using sulfuric acid. After separation of the layers, 0.1 g of active coal, Norit SX Ultra, was added to the toluene solution. The toluene solution was filtered in order to remove the active coal, and subsequently concentrated to 15 ml by evaporation. Cooling to -10°C resulted in crystallization. The crystals were filtered, washed with 5 ml of cold toluene and then dried under vacuum at room temperature. The crystals were analyzed qualitatively by TLC (Merck silicagel 60F, d=0.25 mm, art.nr. 5715; mobile phase chloroform /methanol in a ratio of 30/1), detection by UV at 254 nm (sensitivity is 0.3% at 0.1 mg of run product) and by iodine staining (sensitivity 0.1% at 0.1 mg of run product). Results of these treatments are shown in Table I.

Table 1. Effect of various heat treatment parameters on the purity of lovastatin crystals. Qualitative analysis by TLC with UV-detection (sensitivity 0.3% for 0.1 mg of run product) and iodine staining (sensitivity 0.1% for 0.1 mg of run product)

pH	Temperature / °C	Time / minutes	4-acetyl lovastatin in crystal	
			UV*	iodine*
10	21	30	+	+
10	60	30	+	+
10	90	10	+	+
11	90	10	-	-
11	90	5	-	o
12	21	90	-	+
12	60	30	-	-
12	60	10	-	o
12	90	10	-	-
12	90	5	-	o

* - not detectable

o weak spot

+ detectable

EXAMPLE IV: REACTION OF THE PURE COMPOUND 4-ACETYL LOVASTATIN UPON HEAT TREATMENT IN AQUEOUS SOLUTION AT HIGH pH

(a) Preparation of 4-acetyl lovastatin

Acetic anhydride (7 ml; 0.073 mol) was added in one shot to pure lovastatin (25 g; 0.062 mol) and 4-dimethylamino pyridine (1.53 g; 0.013 mol, 20%) in dry pyridine (120 mL at 0°C under nitrogen. The mixture was stirred at 0°C for 6 hours. By TLC-analysis (see Example 3 for description of the method) of

- 10 -

the reaction mixture it was shown that all lovastatin has disappeared, presumably converted into 4-acetyl lovastatin.

Subsequently the pyridine was removed by evaporation and ethyl acetate was added (240 mL) The solution was washed with
5 240 mL of a saturated solution of NaCl. The layers were separated, and the organic layer was dried over anhydrous magnesium sulfate. The organic layer was filtered and subsequently ethyl acetate was evaporated, yielding a light yellow oil. This oil was identified as a 1/1 mixture of 4-acetyl
10 lovastatin and dehydro lovastatin by proton NMR.

The 4-acetyl lovastatin appeared to be unstable upon storage under N₂ and further conversion of 4-acetyl lovastatin to dehydro lovastatin occurred.

15 (b) Purification of 4-acetyl lovastatin by chromatography

3 g of a 1/2 mixture of 4-acetyl lovastatin and dehydro lovastatin were dissolved in 2 ml of chloroform/methanol (40/1). Subsequently the solution was absorbed onto 120 g of silicagel (Baker 533), which in turn was developed under pressure (0.3
20 bar) with a mixture of chloroform/methanol (ratio 40/1). Four fractions were collected of which the solvent was removed by evaporation. The third fraction contained 4-acetyl lovastatin with a trace of dehydro lovastatin (0.28 g), and the fourth contained only 4-acetyl lovastatin (0.3 g).

25

(c) Reaction of 4-acetyl lovastatin in an aqueous solution at high pH and elevated temperature

0.3 g of 4-acetyl lovastatin (fourth fraction of Example IV b) was dissolved in a mixture of 2 mL of N,N-dimethylformamide (DMF, Merck) and 98 mL of demineralized water.
30 The pH was adjusted to pH 12.5 with NaOH, and the solution was stirred at 60°C for 1 hour.

Subsequently the reaction mixture was cooled to room temperature, then the pH was adjusted to pH=4 with sulfuric acid
35 and 60 mL of toluene were mixed with the aqueous solution for 0.5 hour in order to extract the reaction products.

- 11 -

After separation of the layers, the toluene solution was heated at 90°C for 6 hours. The toluene was then removed by evaporation, yielding a small amount of product. This product was identified by proton NMR as predominantly dehydro
5 lovastatin, while it did not contain any lovastatin.

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

INTERNATIONAL FORM

Gist-brocades N.V.
Research & Development / Stamconservering
Postbus 1
2600 MA DELFT
Nederland

*name and address of the party to whom the
viability statement is issued*

VIABILITY STATEMENT


issued pursuant to Rule 10.2 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified on the following page

I. DEPOSITOR	II. IDENTIFICATION OF THE MICROORGANISM
<p>Name: Gist-brocades N.V. Research & Development / Stamconservering</p> <p>Address: Postbus 1 2600 MA DELFT Nederland</p>	<p>Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: CBS 456.95</p> <p>Date of the deposit or of the transfer:¹ Friday, 2 June 1995</p>
III. VIABILITY STATEMENT	
<p>The viability of the microorganism identified under II above was tested on Tuesday, 13 June 1995². On that date, the said microorganism was</p> <p><input checked="checked" type="checkbox"/>³ viable</p> <p><input type="checkbox"/>³ no longer viable</p>	

¹ Indicate the date of the original deposit or, where a new deposit or a transfer has been made,
the most recent relevant date (date of the new deposit or date of the transfer).

² In the cases referred to in Rule 10.2(a)(ii) and (iii), refer to the most recent viability test.

³ Mark with a cross the applicable box.

IV. CONDITIONS UNDER WHICH THE VIABILITY HAS BEEN PERFORMED ⁴	
V. INTERNATIONAL DEPOSITARY AUTHORITY	
Name: Centraalbureau voor Schimmelcultures	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s):
Address: Oosterstraat 1 P.O. Box 273 3740 AG BAARN The Netherlands	 drs F.M. van Asma dr M.C. Agterberg
	Date: Monday, 3 July 1995

⁴ Fill in if the information has been requested and if the results of the test were negative.


BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

INTERNATIONAL FORM

Gist-brocades N.V.
Research & Development / Stamconservering
Postbus 1
2600 MA DELFT
Nederland

name and address of depositor

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR: DS28373	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: CBS 456.95
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by: <input type="checkbox"/> a scientific description <input checked="" type="checkbox"/> a proposed taxonomic designation (mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary accepts the microorganism identified under I above, which was received by it on Friday, 2 June 1995 (date of the original deposit) ¹	
IV. RECEIPT OF REQUEST FOR CONVERSION	
The microorganism identified under I above was received by this International Depositary Authority on not applicable (date of the original deposit) and a request to convert the original deposit to a deposit under the Budapest Treaty was received by it on not applicable (date of receipt of request for conversion)	
V. INTERNATIONAL DEPOSITARY AUTHORITY	
Name: Centraalbureau voor Schimmelcultures Address: Oosterstraat 1 P.O. Box 273 3740 AG BAARN The Netherlands	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s):  drs F.M. van Asma dr M.C. Agterberg Date: Monday, 3 July 1995

¹ Where Rule 6.4(d) applies, such date is the date on which the status of international
depositary authority was acquired.

5

CLAIMS

1. A process to improve the recovery of lovastatin,
10 pravastatin or compactin from a broth filtrate, comprising:
- growing in a medium microorganisms that produce a member
of the group consisting of lovastatin, compactin or
pravastatin resulting in a product medium
- removal of the biomass from the product medium to obtain
15 a clarified broth filtrate
- isolating the purified lovastatin, compactin or
pravastatin, respectively,
characterized by adjusting the pH of the clarified broth
filtrate above about pH 10.

20

2. A process according to claim 1, wherein the pH is
between pH=10 and pH=13.

3. A process according to claim 2, wherein the pH is
25 between pH=11 and pH=12.5.

4. A process according to any one of the claims 1-3,
characterized by heating the broth filtrate above about 50°C.

30 5. A process according to claim 4, wherein the temperature
is between 60 and 90°C.




Figure 1




Figure 2

INTERNATIONAL SEARCH REPORT

National Application No.

PCT/EP 96/03495

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07C69/732 C07C67/317 C12P7/62

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07C C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	GB,A,2 255 974 (MERCK&CO INC.) 25 November 1992 see page 13 - page 14; claims ---	1
A	US,A,5 159 104 (REBECCA L. DABORA) 27 October 1992 cited in the application see column 2 see column 4, line 41 - column 5, line 6 see column 6; claims 1-3 -----	1

☐

Further documents are listed in the continuation of box C.

☒

Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

4 November 1996

Date of mailing of the international search report

08.11.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (- 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (- 31-70) 340-3016

Authorized officer

Kinzinger, J

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 96/03495

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB-A-2255974	25-11-92	NONE	
US-A-5159104	27-10-92	AT-T- 141584	15-09-96
		CA-A- 2067722	02-11-92
		DE-D- 69212884	26-09-96
		EP-A- 0511867	04-11-92
		ES-T- 2090507	16-10-96
		JP-A- 5230052	07-09-93
		JP-B- 7064837	12-07-95

